EpiPen Auto Injector Directions Twinject ® 0.3mg and 0.15mg Directions First remove the EpiPen Auto Injector Remove caps from the plastic carrying case labeled "1" and Pull of the blue safety release cap "2" Place rounded tip against outer thigh, press down hard until needle penetrates Hold for 10 seconds and remove SECOND DOSE ADMINISTRATION Hold orange tip near outer thigh If symptoms don't improve after 10 mins. (always apply to thigh) administer second dose. Unscrew rounded tip. Pull syringe from Swing and firmly push orange tip against outer thigh. Hold on thigh for barrel by holding the blue collar at the needle base. approximately 10 seconds. Remove the pe the EPIPEN from the thigh and massage area for 10 sec. Slide yellow collar off plunger Adrenaclick ™0.3mg and .15mg Directions Put needle into thigh through Remove Grey cap labeled "1" skin, push plunger down all the way, and remove. Staff Trained To Administer Epi Pen Remove Grey cap 1. _ labeled "2" 2. _____ 3. Place red rounded tip against outer thigh, press down hard until 4. needle penetrates. Hold for 10 sec. and remove 5. 6.

KAUFMAN ISD EPINEPHRINE AUTO INJECTOR USE

Monitoring

Stay with the student; alert healthcare professionals and parent. Tell rescue squad Epi was given; request ambulance with epinephrine. Note time epinephrine was administered. A second dose of epinephrine can be given 5 minutes or more after the first dose if symptoms persist or recur. For severe reaction, consider keeping student lying on back with less raised. Treat student even if parents cannot be reached

with legs faised. Theat student even in parents cannot be reached.	
Student Name	_ Teacher
School Nurse	
CALL 911/Doctor	Phone
Parent/Guardian	Phone